


ViralPolish® Purification (VP) Resin

ViralPolish® chromatography media is specifically for virus purification; Comprising dual layer agarose beads with an inert external shell with tightly controlled pore size and internal polyfunctional ligands for rapid high capacity binding of impurities.



Large particles, such as viruses, are excluded from the beads. The beads can be packed in a column where the virus will pass through the column bed and collected in the flow through fraction, whilst impurities will be captured within the beads. This affords a very gentle purification process resulting in efficient clean up and high recovery of active virus particles.

ViralPolish® Resin is available in three different outer shell pore sizes, and two different impurity binding chemistries (see table below).

1. Properties

ViralPolish® Resin range of chromatography media has a thin porous inert out-layer that excludes molecules with molecular weight >10000 KDa (10000A and 10000B), >9000 KDa (9000A and 9000B), >8000 KDa (8000A and 8000B), >7000 KDa (7000A and 7000B), >6000 KDa (6000A and 6000B), >5000 KDa (5000A and 5000B), >700 KDa (700A and 700B), >400 KDa (400A and 400B), respectively. Molecules smaller than the designated size-exclusion level can penetrate the outer layer and are adsorbed by the mixed-mode ligands in the core.

The base matrix is a composite of polysaccharides that have been highly cross-linked. The media is stable in most of the chemical conditions experienced in the bioprocessing industry.

Selection Guide


Shelled Mixed-Mode Resin	Application Guide	Key Features
ViralPolish® 10000A ViralPolish® 10000B	Suitable for very large viruses and large exosomes; or for removing DNAs > 1kb	<p>The inert out shell has molecular cut-off of 10000kDa, 9000kDa, 8000kDa, 7000kDa, 6000kDa, 5000kDa, 700kDa and 400kDa, respectively, to block viral particles of different sizes.</p> <p>“A”: Mixed-mode anion ligand with mild hydrophobicity; can be easily re-generated and re- used; but its binding capacity may be compromised.</p> <p>“B”: Mixed-mode anion ligand with strong hydrophobicity; have higher loading capacity but is difficult to re-generate. They are more suitable for single-use.</p>
ViralPolish® 9000A ViralPolish® 9000B	Suitable for large viruses and large exosomes; or for removing DNAs > 1kb	
ViralPolish® 8000A ViralPolish® 8000B	Suitable for medium to large size viruses; or for removing DNAs of 1kb	
ViralPolish® 7000A ViralPolish® 7000B	Suitable for small to medium size viruses; or for removing DNAs of 0.5 - 1kb	
ViralPolish® 6000A ViralPolish® 6000B	Suitable for small viruses	
ViralPolish® 5000A ViralPolish® 5000B	<p>Good for lentivirus, adeno virus, VLPs and other viruses of large sizes</p> <p>AAV may be purified if very high molecular-weight impurity (e.g. truncated AAV or empty capsid) need be removed.</p>	
ViralPolish® 700A ViralPolish® 700B	Good for AAVs, other viruses and VLPs of similar sizes, plasmids etc	
ViralPolish® 400A ViralPolish® 400B	Good for certain vaccine antigens, large proteins etc	

Note: other mixed-mode ligands are available on request.

Product characteristics

Particle size	50 – 150 µm
Ligand	Mixed-mode
Ligand density	> 50 µmol / ml resin
Protein binding capacity	Depends on the type of proteins and binding conditions; could be > 40 mg / ml resin
Operational pressure	Up to 3 bar
Flow velocity	Up to 500 cm/hr
Chemical stability	Stable in most common aqueous buffers
pH stability	2-14 (<2 h) 3-12 (long term)
Storage	20% ethanol at 4°C

The pore size distribution of the outer layer is critical to prevent molecules of a given molecular weight from binding to exposed 'active core'. ViralPolish® Resin has tightly controlled pores giving best in class exclusion of virus and large molecules, resulting in significantly higher recovery of active viral particles.



2. Applications

This media is ideal for rapid clean-up, either as a first step or as a second step, to purify virus particles. The beads can also be used in batch mode; when added to a flask of virus containing media the beads will selectively allow small molecules to enter and be retained, whilst the virus is excluded and remains in the external media.

A typical purification strategy for virus would be to use ViralPolish® Resin as an initial step, to clean up the sample, prior to binding and elution, typically from an anion exchange media designed for virus purification.

Alternatively, very dilute samples can be purified by first passing through a Q anion exchange column to concentrate the sample followed by passing through ViralPolish® column to remove residual impurities.

3. Operation

The loose media is stored in 20% ethanol on delivery. It can be easily packed to any commercially available chromatography columns.

Column packing can be done in deionised water or low salt buffers using all the common methods. For flow packing, particular attention should be given to the maximum packing pressure. The typical packing pressure is 0.2 – 0.3 MPa (2-3 bar). Increase or decrease the packing pressure if the peak asymmetry becomes >1.5 or <0.7. Operate the column at a pressure lower than the maximum packing pressure.

Packing Efficiency Assessment

To check the quality of the packing and to monitor this during the working life of the column, column efficiency should be tested directly after packing, prior to re-use and if there is an observed deterioration in separation performance. The efficiency of a packed column is expressed in terms of the height equivalent to a theoretical plate (HETP) and

the asymmetry factor (As). These values are easily determined by applying a sample such as 1% acetone solution to the column and using water as eluent. Sodium chloride can also be used as a test substance. Use a concentration of 2 M NaCl in water with 0.4 M NaCl in water as eluent. It is important that conditions and equipment are kept constant so that results are comparable. Changes in solute, solvent, eluent, sample volume, flow rate, liquid pathway, temperature, etc., will influence the results. A sample volume of less than 2.5% of the column volume and the flow velocity between 15 and 30 cm/h will give the optimal results.

4. Method optimization

We recommend scouting for optimal binding pH, ionic strength and flow velocity (i.e. residence time). We recommend special attention be paid to optimising the flow velocity to balance product yield and product purity.

In general, balancing product recovery against process throughput is the major consideration when optimizing a method. However, for the purification of shear-force sensitive molecules, the operational flow velocity needs to be optimised to minimise possible damage to the target molecule.

Tips:

- If unexpectedly high loss of product is noticed, consider using increased flow rates and/or increased ionic strength, or to adjust the pH to lower the charge of the target product.
- If too high level of impurities remain in flow-through mode, another medium with higher size-exclusion level may be tested at increased flow velocity. For example, ViralPolish® 5000 may be used instead of ViralPolish® 700.
- For AAV viruses, the first choice is ViralPolish® 700. For larger viruses, ViralPolish® 5000 is the first choice.

5. Process scale-up

ViralPolish® range of media is designed for bioprocessing use with full regulatory support documents. Please contact us for further information.

6. Maintenance

Depending on individual applications, please see the following recommendations.

Note: when sodium hydroxide solution or organic solvent (e.g. 20% ethanol etc) is used, the flowrate must be less than 50% of the normal operational flowrate, because the column pressure will increase under these chemical conditions.

Cleaning-in-place (CIP)

CIP is a procedure that removes strongly bound materials such as lipids, endotoxins and denatured proteins that remain in the column after regeneration. Regular CIP prevents the build up of contaminants in the packed bed and helps to maintain the column performance.

A specific CIP protocol should be developed for each process according to the type of contaminants present. The frequency of CIP depends on the nature of individual applications.

The following information works as a general guidance.

The contaminants bound hydrophobically can be removed by using the following reagents: 1 M NaOH, low percentage non-ionic detergents (e.g. 0.1 – 2%), 30% isopropanol in basic or acidic conditions (e.g. in the presence of acetic acid or phosphoric acid). A combination of the above reagents can be explored as well. In general, the incubation time should be longer (e.g. from 30 minutes to 2 hours) to ensure full dissociation of the contaminants.

Note: Long contact times should be avoided when using alcohols in acrylic columns.

Sanitization

Sanitization using 0.5-1.0 M NaOH with a contact time of 30 mins is recommended.

7. Storage

The loose media or column should be stored in 20% ethanol to prevent microbial growth. Store the column at a temperature of +2°C to +8°C. After storage, equilibrate the column with at least 5 bed volumes of running buffer before use.

Ordering Information

Note: other column sizes available on request

Product Description	Pack Size	Product Code
ViralPolish® Virus Purification Resin, 400A	25ml	VP-400A-25
	100ml	VP-400A-100
	1L	VP-400A-1L
	5L	VP-400A-5L
	10L	VP-400A-10L
ViralPolish® Virus Purification Resin, 400B	25ml	VP-400B-25
	100ml	VP-400B-100
	1L	VP-400B-1L
	5L	VP-400B-5L
	10L	VP-400B-10L
ViralPolish® Virus Purification Resin, 700A	25ml	VP-700A-25
	100ml	VP-700A-100
	1L	VP-700A-1L
	5L	VP-700A-5L
	10L	VP-700A-10L
ViralPolish® Virus Purification Resin, 700B	25ml	VP-700B-25
	100ml	VP-700B-100
	1L	VP-700B-1L
	5L	VP-700B-5L
	10L	VP-700B-10L
ViralPolish® Virus Purification Resin, 5000A	25ml	VP-5000A-25
	100ml	VP-5000A-100
	1L	VP-5000A-1L
	5L	VP-5000A-5L
	10L	VP-5000A-10L
ViralPolish® Virus Purification Resin, 5000B	25ml	VP-5000B-25
	100ml	VP-5000B-100
	1L	VP-5000B-1L
	5L	VP-5000B-5L
	10L	VP-5000B-10L
ViralPolish® Virus Purification Resin, 6000B	25ml	VP-6000B-25
	100ml	VP-6000B-100
	1L	VP-6000B-1L
	5L	VP-6000B-5L
	10L	VP-6000B-10L

ViralPolish® Virus Purification Resin, 7000B	25ml 100ml 1L 5L 10L	VP-7000B-25 VP-7000B-100 VP-7000B-1L VP-7000B-5L VP-7000B-10L
ViralPolish® Virus Purification Resin, 8000B	25ml 100ml 1L 5L 10L	VP-8000B-25 VP-8000B-100 VP-8000B-1L VP-8000B-5L VP-8000B-10L
ViralPolish® Virus Purification Resin, 9000B	25ml 100ml 1L 5L 10L	VP-9000B-25 VP-9000B-100 VP-9000B-1L VP-9000B-5L VP-9000B-10L
ViralPolish® Virus Purification Resin, 10000B	25ml 100ml 1L 5L 10L	VP-10000B-25 VP-10000B-100 VP-10000B-1L VP-10000B-5L VP-10000B-10L



A CALIBRE SCIENTIFIC COMPANY

Calibre Scientific UK
Unit 5A, R-Evolution @
The Advanced Manufacturing Park
Selden Way
Rotherham
S60 5XA

Tel: +44 (0) 114 224 2257
Email: info@proteinark.com
Web: www.proteinark.com

All goods and services are sold subject to the terms and conditions of sale of Protein Ark Ltd. The company reserves the rights, subject to regulatory or contractual approval, if required, to make changes in the specifications and features shown herein, or discontinue the products described at any time without notice or obligation. Contact Protein Ark Ltd for the most current information.

